

# Package ‘HIVcDNAvantWout03’

October 17, 2024

**Version** 1.44.0

**Date** 2005-01-31

**Author** Dr. Angelique van't Wout, Department of Microbiology,  
University of Washington

**Title** T cell line infections with HIV-1 LAI (BRU)

**Description** The expression levels of approximately 4600 cellular RNA transcripts were assessed in CD4+ T cell lines at different times after infection with HIV-1BRU using DNA microarrays. This data corresponds to the first block of a 12 block array image (001030\_08\_1.GEL) in the first data set (2000095918 A) in the first experiment (CEM LAI vs HI-LAI 24hr). There are two data sets, which are part of a dye-swap experiment with replicates, representing the Cy3 (green) absorption intensities for channel 1 (hiv1raw) and the Cy5 (red) absorption intensities for channel 2 (hiv2raw).

**biocViews** ExperimentData, MicroarrayData, TwoChannelData, HIVData

**License** GPL (>= 2)

**Maintainer** Chris Fraley <fraley@stat.washington.edu>

**URL** <http://expression.microslu.washington.edu/expression/vantwoutjvi2002.html>

**git\_url** <https://git.bioconductor.org/packages/HIVcDNAvantWout03>

**git\_branch** RELEASE\_3\_19

**git\_last\_commit** 8751bcf

**git\_last\_commit\_date** 2024-04-30

**Repository** Bioconductor 3.19

**Date/Publication** 2024-10-17

## Contents

hiv1raw	2
hiv2raw	3
<b>Index</b>	<b>4</b>

---

hiv1raw

*T cell line infections with HIV-1 LAI (BRU)*

---

### Description

The expression levels of approximately 4600 cellular RNA transcripts were assessed in CD4+ T cell lines at different times after infection with HIV-1BRU using DNA microarrays. There are two data sets, which are part of a dye-swap experiment with replicates, representing the Cy3 (green) absorption intensities for channel 1 (hiv1raw) and the Cy5 (red) absorption intensities for channel 2 (hiv2raw).

### Usage

```
data(hiv1raw)
```

### Format

This data represents a block within a microarray image with 12x32 spots. It is stored as a vector of length 450,000 representing a 450x1000 matrix (ordered by column) of intensities encoded for compact (16-bit TIFF) storage.

### Details

The intensities can be obtained from this data by first subtracting them from 65535, then squaring, then multiplying by a scale factor 4.71542407E-05. In other words, a number  $x$  in the hiv1 data set corresponds to intensity  $(256 * 256 - 1 - x)^2 * .0000471542407$ .

### Source

Dr. Angelique van't Wout, Department of Microbiology, University of Washington

The data corresponds to the first block of a 12 block array image ('001030\08\1.GEL') in the first data set ('2000095918 A') in the first experiment ('CEM LAI vs HI-LAI 24hr') of the following data archive: <http://expression.microslu.washington.edu/expression/vantwoutjvi2002.html>

### References

van't Wout AB, Lehrman GK, Mikheeva SA, O'Keeffe GC, Katze MG, Bumgarner RE, Geiss GK and Mullins JI, Cellular gene expression upon human immunodeficiency virus type 1 infection of CD4(+)-T-cell lines, *J Virol.* 2003 Jan;77(2):1392-402.a

---

hiv2raw

*T cell line infections with HIV-1 LAI (BRU)*

---

### Description

The expression levels of approximately 4600 cellular RNA transcripts were assessed in CD4+ T cell lines at different times after infection with HIV-1BRU using DNA microarrays. There are two data sets, which are part of a dye-swap experiment with replicates, representing the Cy3 (green) absorption intensities for channel 1 (hiv1raw) and the Cy5 (red) absorption intensities for channel 2 (hiv2raw).

### Usage

```
data(hiv2raw)
```

### Format

This data represents a block within a microarray image with 12x32 spots. It is stored as a vector of length 450,000 representing a 450x1000 matrix (ordered by column) of intensities encoded for compact (16-bit TIFF) storage.

### Details

The intensities can be obtained from this data by first subtracting them from 65535, then squaring, then multiplying by a scale factor 4.71542407E-05. In other words, a number  $x$  in the hiv1 data set corresponds to intensity  $(256 * 256 - 1 - x)^2 * .0000471542407$ .

### Source

Dr. Angelique van't Wout, Department of Microbiology, University of Washington

The data corresponds to the first block of a 12 block array image ('001030\08\1.GEL') in the first data set ('2000095918 A') in the first experiment ('CEM LAI vs HI-LAI 24hr') of the following data archive:\a href="http://expression.microslu.washington.edu/expression/vantwoutjvi2002.html">http://expression.microslu.washington.edu/expression/vantwoutjvi2002.html

### References

van't Wout AB, Lehrman GK, Mikheeva SA, O'Keeffe GC, Katze MG, Bumgarner RE, Geiss GK and Mullins JI, Cellular gene expression upon human immunodeficiency virus type 1 infection of CD4(+)-T-cell lines, *J Virol.* 2003 Jan;77(2):1392-402.

# Index

\* **datasets**

hiv1raw, [2](#)

hiv2raw, [3](#)

hiv1raw, [2](#)

hiv2raw, [3](#)