

# *vtpNet*: variant-transcription factor-phenotype networks

VJ Carey

April 30, 2026

## 1 Introduction

In a wide-ranging paper (PMID 22955828 Maurano et al. (2012)), Maurano and colleagues illustrate the concept of “common networks for common diseases” with a bipartite graph. One class of nodes is a set of autoimmune disorders, the other class is a set of transcription factors (TFs). In this graph, an edge exists between a disorder node and a TF node if a SNP that is significantly associated with the risk of the disorder lies in a genomic region possessing a strong match to the binding motif of the TF. This package defines tools to investigate the construction and statistical interpretation of such bipartite graphs, which we will denote VTP (variant-transcription factor-phenotype) networks.

## 2 Illustrative example of an unpruned VTP

The following code uses the `graphNEL` class to construct an approximation to the complete bipartite graph underlying Figure 4A of the Maurano paper; Figure 1 illustrates an arbitrary complete subgraph. The elements of `diseaseTags` are formatted to allow multiline rendering of the strings in node displays. It will be useful to distinguish a display token type and an analysis token type to simplify programming.

```
> #  
> # tags formatted for display  
> #  
> diseaseTags = c("Ankylosing\\nspondylitis", "Asthma",  
+ "Celiac\\ndisease", "Crohn's\\ndisease",  
+ "Multiple\\nsclerosis", "Primary\\nbiliary\\ncirrhosis",  
+ "Psoriasis", "Rheumatoid\\narthrititis",  
+ "Systemic\\nlupus\\nerythematosus",  
+ "Systemic\\nsclerosis", "Type 1\\ndiabetes",
```

```

+       "Ulcerative\\nocolitis"
+ )
> TFtags = c("ELF3", "MEF2A", "TCF3", "PAX4", "STAT3",
+   "ESR1", "POU2F1", "STAT1", "YY1", "SP1", "CDC5L",
+   "NR3C1", "EGR1", "PPARG", "HNF4A", "REST", "PPARA",
+   "AR", "NFKB1", "HNF1A", "TFAP2A")
> # define adjacency matrix
> adjm = matrix(1, nr=length(diseaseTags), nc=length(TFtags))
> dimnames(adjm) = list(diseaseTags, TFtags)
> library(graph)
> cvtp = ugraph(aM2bpG(adjm)) # complete (V)TP network; variants not involved yet

```

### 3 Data on GWAS variants: their associated phenotype, locations, and other characteristics

We will use the GWAS data provided at <https://www.sciencemag.org/content/suppl/2012/09/04/science.1222794.DC1/1222794-Maurano-tableS2.txt>, which was manually imported to a GRanges instance in hg19 origin-1 coordinates.

```

> library(vtpnet)
> data(maurGWAS)
> length(maurGWAS)

[1] 5654

> names(values(maurGWAS))

[1] "name"                "disease_trait"
[3] "disease_class"       "internally_replicated"
[5] "independently_replicated" "In_DHS"
[7] "fetal_origin"        "X.LOG.P."
[9] "sample_size"

```

### 4 Data on transcription factor binding sites

We have included the result of using FIMO Grant et al. (2011) to scan for motif matches for TF PAX4 as modeled in the Bioconductor *MotifDb* collection. The `-max-stored-scores` parameter was set to 10000000 so that  $p$  of up to  $10^{-4}$  are retained.

```

> data(pax4)
> length(pax4)

```

```

> library(Rgraphviz)
> #flat = function(x, g) c(x, edges(g)[[x]])
> #sub = subGraph(unique(c(flat("Crohn's\\ndisease", cvtp),
> #   flat("Ulcerative\\ncolitis", cvtp))), cvtp)
> sub = subGraph(unique(c(diseaseTags[1:4], TFtags[1:6])), cvtp)
> plot(sub, attrs=list(node=list(shape="box", fixedsize=FALSE)))
> #plot(cvtp, attrs=list(graph=list(margin=c(.5,.5), size=c(4.1,4.1)),
> #   node=list(shape="box", fixedsize=FALSE, height=1)))

```




Figure 1: A complete bipartite graph for arbitrarily selected subsets of the autoimmune disorders and TFs found in Figure 4A of Maurano et al.

```
[1] 1862156
```

```
> pax4[1:4]
```

GRanges object with 4 ranges and 8 metadata columns:

|     | seqnames<br><Rle> | ranges<br><IRanges> | strand<br><Rle> | source<br><factor> | type<br><factor> | score<br><numeric> |
|-----|-------------------|---------------------|-----------------|--------------------|------------------|--------------------|
| [1] | chr1              | 10273-10302         | +               | fimo               | nucleotide_motif | 999.917            |
| [2] | chr1              | 10279-10308         | +               | fimo               | nucleotide_motif | 999.962            |
| [3] | chr1              | 11703-11732         | -               | fimo               | nucleotide_motif | 999.999            |
| [4] | chr1              | 11704-11733         | -               | fimo               | nucleotide_motif | 999.955            |

|     | phase<br><integer> | Name<br><character>    | pvalue<br><character> | qvalue<br><character> |
|-----|--------------------|------------------------|-----------------------|-----------------------|
| [1] | <NA>               | +Mmusculus-JASPAR_CO.. | 8.35e-05              | 0.396                 |
| [2] | <NA>               | +Mmusculus-JASPAR_CO.. | 3.79e-05              | 0.361                 |
| [3] | <NA>               | -Mmusculus-JASPAR_CO.. | 8.04e-07              | 0.194                 |
| [4] | <NA>               | -Mmusculus-JASPAR_CO.. | 4.46e-05              | 0.368                 |

|     | sequence<br><character> |
|-----|-------------------------|
| [1] | TAACCCTAACCCCTAACCCCA.. |
| [2] | TAACCCTAACCCCAACCCCA..  |
| [3] | AAAAAAAATACACATGGCCAG.. |
| [4] | TAAAAAAAATACACATGGCCA.. |

-----  
seqinfo: 92 sequences from an unspecified genome; no seqlengths

We can also generate our own motif-match ranges. Here is an example of a parallelized search against hg19 using matchPWM.

```
> library(foreach)
> library(doParallel)
> registerDoParallel(cores=12)
> library(BSgenome.Hsapiens.UCSC.hg19)
> library(MotifDb)
> sn = seqnames(Hsapiens)[1:24]
> pax4 = query(MotifDb, "pax4")[[1]]
> ans = foreach(i=1:24) %dopar% {
+   cat(i)
+   subj = Hsapiens[[ sn[i] ]]
+   matchPWM( pax4, subj, "75%" )
+ }
> pax4_75 =
+ do.call(c, lapply(1:length(ans), function(x)
```

```
+ {GRanges(sn[x], as(ans[[x]], "IRanges"))})
> save(pax4_75, file="pax4_75.rda")
```

Results of such searches retaining matches at scores of 85% and 75% of the maximum achievable score have been stored with this package.

## 5 Building a VTP network: one edge per phenotype

### 5.1 Raw matches

We can survey the entire GWAS catalog for intersection with putative PAX4 binding sites. First the two Bioconductor internal binding site sets.

```
> data(pax4_85)
> vp_pax4_85 = maurGWAS[ overlapsAny(maurGWAS, pax4_85) ]
> length(vp_pax4_85)
```

```
[1] 0
```

```
> data(pax4_75)
> vp_pax4_75 = maurGWAS[ overlapsAny(maurGWAS, pax4_75) ]
> length(vp_pax4_75)
```

```
[1] 54
```

Then the FIMO-based set.

```
> vp_pax4_fimo = maurGWAS[ overlapsAny(maurGWAS, pax4) ]
> length(vp_pax4_fimo)
```

```
[1] 67
```

The lengths reported here are the numbers of phenotypes linked to PAX4 in a VTP according to various motif matching schemes. For the two non-null results, we have

```
> u75 = unique(vp_pax4_75$disease_trait)
> ufimo = unique(vp_pax4_fimo$disease_trait)
> length(setdiff(u75, ufimo))
```

```
[1] 23
```

```
> length(setdiff(ufimo, u75))
```

```
[1] 28
```

Clearly the identification of TP links is sensitive to the approach used to locate binding sites. However, as noted in the Maurano paper, the use of matching to the reference genome without SNP injection is potentially problematic.

## 5.2 Filtering

It is useful to restrict the phenotypes of interest, and to map them to broader classes, and to include TFBS matching scores for the purpose of filtering edges. Here we will use the NHGRI GWAS catalog against FIMO-based (reference genome matching only) PAX4 calls.

```
> data(cancerMap)
> requireNamespace("gwascat")
> load(system.file("legacy/gwrngs19.rda", package="gwascat"))
> cangw = filterGWASbyMap( gwrngs19, cancerMap )
> getOneHits( pax4, cangw, "fimo" )
```

GRanges object with 8 ranges and 41 metadata columns:

|       | seqnames                                      | ranges                 | strand                 | Date.Added.to.Catalog | PUBMEDID  |
|-------|---|------------------------|------------------------|-----------------------|-----------|
|       | <Rle>   | <IRanges>              | <Rle>                  | <character>           | <integer> |
| 3475  | chr8  | 129194641              | *                      | 09/12/2013            | 23535729  |
| 3480  | chr11   | 65583066               | *                      | 09/12/2013            | 23535729  |
| 6963  | chr2  | 26526419               | *                      | 01/25/2013            | 23144319  |
| 7155  | chr6  | 143943314              | *                      | 01/15/2013            | 23108145  |
| 7480  | chr20   | 32588095               | *                      | 11/30/2012            | 22976474  |
| 12585 | chrX  | 37854727               | *                      | 11/15/2010            | 20932654  |
| 13650 | chr12   | 14653867               | *                      | 07/12/2010            | 20543847  |
| 15145 | chr10   | 63752159               | *                      | 09/04/2009            | 19684604  |
|       | First.Author                                  | Date                   | Journal                |                       |           |
|       | <character>                                   | <character>            | <character>            |                       |           |
| 3475  | Michailidou K                                 | 04/01/2013             | Nat Genet              |                       |           |
| 3480  | Michailidou K                                 | 04/01/2013             | Nat Genet              |                       |           |
| 6963  | Lee Y   | 11/08/2012             | Carcinogenesis         |                       |           |
| 7155  | Wang LE                                       | 10/29/2012             | Cancer Res             |                       |           |
| 7480  | Siddiq A                                      | 09/13/2012             | Hum Mol Genet          |                       |           |
| 12585 | Kerns SL                                      | 10/05/2010             | Int J Radiat Oncol B.. |                       |           |
| 13650 | Turnbull C                                    | 06/13/2010             | Nat Genet              |                       |           |
| 15145 | Papaemmanuil E                                | 08/16/2009             | Nat Genet              |                       |           |
|       | Link  | Study                  | Disease.Trait          |                       |           |
|       | <character>                                   | <character>            | <character>            |                       |           |
| 3475  | http://www.ncbi.nlm... Large-scale genotypi.. |                        | Breast cancer          |                       |           |
| 3480  | http://www.ncbi.nlm... Large-scale genotypi.. |                        | Breast cancer          |                       |           |
| 6963  | http://www.ncbi.nlm... Prognostic implicati.. | Non-small cell lung .. |                        |                       |           |
| 7155  | http://www.ncbi.nlm... Genome-wide associat.. | Lung Cancer (DNA rep.. |                        |                       |           |
| 7480  | http://www.ncbi.nlm... A meta-analysis of g.. |                        | Breast cancer          |                       |           |
| 12585 | http://www.ncbi.nlm... Genome-wide associat.. | Erectile dysfunction.. |                        |                       |           |
| 13650 | http://www.ncbi.nlm... Variants near DMRT1,.. | Testicular germ cell.. |                        |                       |           |

|       |  |                           |                     |                        |
|-------|--|---------------------------|---------------------|------------------------|
| 15145 | http://www.ncbi.nlm... Loci on 7p12.2, 10q2.. Acute lymphoblastic .. |                           |                     |                        |
|       | Initial.Sample.Size  | Replication.Sample.Size   | Region              | Chr_id                 |
|       | <character>  | <character>               | <character>         | <character>            |
| 3475  | 10,052   | European ance..           | 45,290              | European ance..        |
| 3480  | 10,052   | European ance..           | 45,290              | European ance..        |
| 6963  | 348  | Korean ancestry ..        | NR                  | 2p23.3                 |
| 7155  | 914  | European ancestr..        | 679                 | European ancestr..     |
| 7480  | 3,666  | European ances..          | 562                 | European ancestr..     |
| 12585 | 27   | African American ..       | <NA>                | Xp11.4                 |
| 13650 | 979  | European ancestr..        | 664                 | European ancestr..     |
| 15145 | 907  | European ancestr..        | <NA>                | 10q21.2                |
|       | Chr_pos.hg38   | Reported.Gene.s.          | Mapped_gene         |                        |
|       | <numeric>  | <character>               | <character>         |                        |
| 3475  | 128182395  | MIR1208, MYC              | MIR1208 - LINC01263 |                        |
| 3480  | 65815595   | DKFZp761E198, OVOL1,..    | OVOL1-AS1 - SNX32   |                        |
| 6963  | 26303551   | GPR113                    | HADHB - GPR113      |                        |
| 7155  | 143622177  | PHACTR2                   | PHACTR2             |                        |
| 7480  | 34000289   | RALY, EIF2S2, ASIP        | RALY                |                        |
| 12585 | 37995474   | SYTL5                     | CXorf27 - SYTL5     |                        |
| 13650 | 14500933   | ATF7IP                    | ATF7IP              |                        |
| 15145 | 61992400   | ARID5B                    | ARID5B              |                        |
|       | Upstream_gene_id   | Downstream_gene_id        | Snps_gene_ids       | Upstream_gene_distance |
|       | <character>  | <character>               | <character>         | <character>            |
| 3475  | 100302281  | 101927774                 |                     | 32.21                  |
| 3480  | 101927828  | 254122                    |                     | 24.73                  |
| 6963  | 3032   | 165082                    |                     | 13.09                  |
| 7155  | <NA>   | <NA>                      | 9749                | <NA>                   |
| 7480  | <NA>   | <NA>                      | 22913               | <NA>                   |
| 12585 | 25763  | 94122                     |                     | 4.16                   |
| 13650 | <NA>   | <NA>                      | 55729               | <NA>                   |
| 15145 | <NA>   | <NA>                      | 84159               | <NA>                   |
|       | Downstream_gene_distance   | Strongest.SNP.Risk.Allele | SNPs                |                        |
|       | <character>  | <character>               | <character>         |                        |
| 3475  | 222.87   | rs11780156-T              | rs11780156          |                        |
| 3480  | 18.24  | rs3903072-G               | rs3903072           |                        |
| 6963  | 4.62   | rs6753473-G               | rs6753473           |                        |
| 7155  | <NA>   | rs9390123-A               | rs9390123           |                        |
| 7480  | <NA>   | rs2284378-T               | rs2284378           |                        |
| 12585 | 11.11  | rs872690-?                | rs872690            |                        |
| 13650 | <NA>   | rs2900333-C               | rs2900333           |                        |
| 15145 | <NA>   | rs7089424-C               | rs7089424           |                        |
|       | Merged Snp_id_current  | Context                   | Intergenic          |                        |

|       | <character> | <character> | <character> | <character> |
|-------|-------------|-------------|-------------|-------------|
| 3475  | 0           | 11780156    | Intergenic  | 1           |
| 3480  | 0           | 3903072     | Intergenic  | 1           |
| 6963  | 0           | 6753473     | Intergenic  | 1           |
| 7155  | 0           | 9390123     | intron      | 0           |
| 7480  | 0           | 2284378     | intron      | 0           |
| 12585 | 0           | 872690      | Intergenic  | 1           |
| 13650 | 0           | 2900333     | UTR-3       | 0           |
| 15145 | 0           | 7089424     | intron      | 0           |

|       | Risk Allele Frequency | p.Value   | Pvalue_mlog | p.Value..text.   | OR.or.beta |
|-------|-----------------------|-----------|-------------|------------------|------------|
|       | <character>           | <numeric> | <numeric>   | <character>      | <numeric>  |
| 3475  | 0.16                  | 3e-11     | 10.52288    |                  | 1.07       |
| 3480  | 0.53                  | 9e-12     | 11.04576    |                  | 1.05       |
| 6963  | 0.052                 | 4e-06     | 5.39794     | (Additive model) | NA         |
| 7155  | 0.3957                | 7e-06     | 5.15490     |                  | NA         |
| 7480  | 0.31                  | 1e-08     | 8.00000     |                  | 1.16       |
| 12585 | 0.03                  | 9e-06     | 5.04576     |                  | 11.78      |
| 13650 | 0.62                  | 6e-10     | 9.22185     |                  | 1.27       |
| 15145 | 0.34                  | 7e-19     | 18.15490    |                  | 1.65       |

|       | X95..CI..text. | Platform..SNPs.passing.QC. | CNV         |
|-------|----------------|----------------------------|-------------|
|       | <character>    | <character>                | <character> |
| 3475  | [1.04-1.10]    | Illumina & Affymetri..     | N           |
| 3480  | [1.04-1.08]    | Illumina & Affymetri..     | N           |
| 6963  | NR             | Affymetrix [271,817]       | N           |
| 7155  | NR             | Illumina [303,669]         | N           |
| 7480  | [1.10-1.22]    | Illumina [2,608,509]..     | N           |
| 12585 | [NR]           | Affymetrix [512,497]       | N           |
| 13650 | [1.12-1.44]    | Illumina [298,782]         | N           |
| 15145 | [1.54-1.76]    | Illumina [291,473]         | N           |

|       | num.Risk Allele Frequency | dclass      | score     | tfstart   | tfend     |
|-------|---------------------------|-------------|-----------|-----------|-----------|
|       | <numeric>                 | <character> | <numeric> | <integer> | <integer> |
| 3475  | 0.1600                    | Breast      | 999.985   | 129194621 | 129194650 |
| 3480  | 0.5300                    | Breast      | 999.952   | 65583065  | 65583094  |
| 6963  | 0.0520                    | Lung        | 999.987   | 26526415  | 26526444  |
| 7155  | 0.3957                    | Lung        | 999.939   | 143943292 | 143943321 |
| 7480  | 0.3100                    | Breast      | 999.928   | 32588075  | 32588104  |
| 12585 | 0.0300                    | Prostate    | 999.903   | 37854721  | 37854750  |
| 13650 | 0.6200                    | Testicular  | 999.990   | 14653848  | 14653877  |
| 15145 | 0.3400                    | ALL (ped)   | 999.962   | 63752142  | 63752171  |

|      | pvalue    | qvalue    |
|------|-----------|-----------|
|      | <numeric> | <numeric> |
| 3475 | 1.49e-05  | 0.318     |

|       |          |       |
|-------|----------|-------|
| 3480  | 4.83e-05 | 0.373 |
| 6963  | 1.25e-05 | 0.310 |
| 7155  | 6.13e-05 | 0.383 |
| 7480  | 7.16e-05 | 0.388 |
| 12585 | 9.72e-05 | 0.403 |
| 13650 | 1.05e-05 | 0.301 |
| 15145 | 3.79e-05 | 0.361 |

-----

seqinfo: 23 sequences from hg19 genome

## 6 Appendix: generating the ALT-injected genome image

```
> altize = function(htag = "21",
+ #
+ # from sketch by Herve Pages, May 2013
+ #
+ slpack="SNPlocs.Hsapiens.dbSNP.20120608",
+ hgpacK = "BSgenome.Hsapiens.UCSC.hg19",
+ faElFun = function(x) sub("%%TAG%%", x, "alt%%TAG%%chr"),
+ faTargFun = function(x)
+   sub("%%TAG%%", x, "alt%%TAG%%_hg19.fa")) {
+   require(slpacK, character.only=TRUE)
+   require(hgpacK, character.only=TRUE)
+   require("ShortRead", character.only=TRUE)
+   chk = grep("ch|chr", htag)
+   if (length(chk)>0) {
+     warning("clearing prefix ch or chr from htag")
+     htag = gsub("ch|chr", "", htag)
+   }
+   snpgettag = paste0("ch", htag)
+   ggettag = paste0("chr", htag)
+   cursnps = getSNPlocs(snpgettag, as.GRanges=TRUE)
+   curgenome = unmasked(Hsapiens[[ggettag]])
+   ref_allele =
+     strsplit(as.character(curgenome[start(cursnps)]),
+       NULL, fixed=TRUE)[[1L]]
+   all_alleles = IUPAC_CODE_MAP[cursnps$alleles_as_ambig]
+   alt_alleles = mapply( function(ref,all)
+     sub(ref, "", all, fixed=TRUE),
+     ref_allele, all_alleles, USE.NAMES=FALSE)
```

```

+     cursnps$ref_allele = ref_allele
+     cursnps$alt_alleles = alt_alleles
+     cursnps$one_alt = substr(cursnps$alt_alleles, 1, 1)
+     altg = list(replaceLetterAt(curgenome, start(cursnps),
+       cursnps$one_alt))
+     names(altg) = faElFun(htag)
+     writeFasta(DNAStringSet(altg), file=faTargFun(htag))
+ }

```

## 7 Session information

```
> sessionInfo()
```

```

R version 4.6.0 (2026-04-24)
Platform: x86_64-pc-linux-gnu
Running under: Ubuntu 24.04.4 LTS

```

```
Matrix products: default
```

```
BLAS: /usr/lib/x86_64-linux-gnu/openblas-pthread/libblas.so.3
```

```
LAPACK: /usr/lib/x86_64-linux-gnu/openblas-pthread/libopenblas-p0.3.26.so; LAPACK ver
```

```
locale:
```

```

[1] LC_CTYPE=en_US.UTF-8      LC_NUMERIC=C
[3] LC_TIME=en_US.UTF-8      LC_COLLATE=en_US.UTF-8
[5] LC_MONETARY=en_US.UTF-8  LC_MESSAGES=en_US.UTF-8
[7] LC_PAPER=en_US.UTF-8     LC_NAME=C
[9] LC_ADDRESS=C             LC_TELEPHONE=C
[11] LC_MEASUREMENT=en_US.UTF-8 LC_IDENTIFICATION=C

```

```
time zone: Etc/UTC
```

```
tzcode source: system (glibc)
```

```
attached base packages:
```

```

[1] parallel stats4 grid stats graphics grDevices utils
[8] datasets methods base

```

```
other attached packages:
```

```

[1] vtpnet_0.52.0      doParallel_1.0.17  iterators_1.0.14
[4] foreach_1.5.2      gwascat_2.44.0     GenomicRanges_1.64.0
[7] Seqinfo_1.2.0      IRanges_2.46.0     S4Vectors_0.50.0
[10] Rgraphviz_2.56.0   graph_1.90.0       BiocGenerics_0.58.0
[13] generics_0.1.4

```

loaded via a namespace (and not attached):

|                               |                             |
|-------------------------------|-----------------------------|
| [1] tidyselect_1.2.1          | dplyr_1.2.1                 |
| [3] blob_1.3.0                | filelock_1.0.3              |
| [5] Biostrings_2.80.0         | bitops_1.0-9                |
| [7] fastmap_1.2.0             | RCurl_1.98-1.18             |
| [9] BiocFileCache_3.2.0       | VariantAnnotation_1.58.0    |
| [11] GenomicAlignments_1.48.0 | XML_3.99-0.23               |
| [13] lifecycle_1.0.5          | survival_3.8-6              |
| [15] KEGGREST_1.52.0          | RSQLite_2.4.6               |
| [17] magrittr_2.0.5           | compiler_4.6.0              |
| [19] rlang_1.2.0              | tools_4.6.0                 |
| [21] yaml_2.3.12              | data.table_1.18.2.1         |
| [23] rtracklayer_1.72.0       | knitr_1.51                  |
| [25] S4Arrays_1.12.0          | bit_4.6.0                   |
| [27] curl_7.1.0               | DelayedArray_0.38.1         |
| [29] abind_1.4-8              | BiocParallel_1.46.0         |
| [31] sys_3.4.3                | SummarizedExperiment_1.42.0 |
| [33] cli_3.6.6                | crayon_1.5.3                |
| [35] httr_1.4.8               | rjson_0.2.23                |
| [37] DBI_1.3.0                | cachem_1.1.0                |
| [39] splines_4.6.0            | AnnotationDbi_1.74.0        |
| [41] XVector_0.52.0           | restfulr_0.0.16             |
| [43] matrixStats_1.5.0        | vctrs_0.7.3                 |
| [45] Matrix_1.7-5             | jsonlite_2.0.0              |
| [47] bit64_4.8.0              | GenomicFeatures_1.64.0      |
| [49] maketools_1.3.2          | snpStats_1.62.0             |
| [51] glue_1.8.1               | codetools_0.2-20            |
| [53] GenomeInfoDb_1.48.0      | BiocIO_1.22.0               |
| [55] UCSC.utils_1.8.0         | tibble_3.3.1                |
| [57] pillar_1.11.1            | rappdirs_0.3.4              |
| [59] BSgenome_1.80.0          | R6_2.6.1                    |
| [61] dbplyr_2.5.2             | httr2_1.2.2                 |
| [63] evaluate_1.0.5           | lattice_0.22-9              |
| [65] Biobase_2.72.0           | png_0.1-9                   |
| [67] Rsamtools_2.28.0         | cigarillo_1.2.0             |
| [69] memoise_2.0.1            | SparseArray_1.12.0          |
| [71] xfun_0.57                | MatrixGenerics_1.24.0       |
| [73] buildtools_1.0.0         | pkgconfig_2.0.3             |

## 8 Bibliography

### References

- Charles E Grant, Timothy L Bailey, and William Stafford Noble. Fimo: scanning for occurrences of a given motif. *Bioinformatics (Oxford, England)*, 27(7):1017–8, Apr 2011. doi: 10.1093/bioinformatics/btr064.
- Matthew T Maurano, Richard Humbert, Eric Rynes, Robert E Thurman, Eric Haugen, Hao Wang, Alex P Reynolds, Richard Sandstrom, Hongzhu Qu, Jennifer Brody, Anthony Shafer, Fidencio Neri, Kristen Lee, Tanya Kutayavin, Sandra Stehling-Sun, Audra K Johnson, Theresa K Canfield, Erika Giste, Morgan Diegel, Daniel Bates, R Scott Hansen, Shane Neph, Peter J Sabo, Shelly Heimfeld, Antony Raubitschek, Steven Ziegler, Chris Cotsapas, Nona Sotoodehnia, Ian Glass, Shamil R Sunyaev, Rajinder Kaul, and John A Stamatoyannopoulos. Systematic localization of common disease-associated variation in regulatory dna. *Science*, 337(6099):1190–5, Sep 2012. doi: 10.1126/science.1222794.